

Homework 12

Physics 2130

Modern Physics

Due: Wednesday December 4, 2002

1. T&Z, #13.15.

(a)

$$\begin{aligned}
 \# \text{ of } ^{14}\text{C} \text{ atoms in 7 g of C} &= N_0 = (1.3 \times 10^{-12}) \frac{7 \text{ g}}{12 \text{ g/mole}} (6.02 \times 10^{23} \text{ atoms/mole}) \\
 &= 4.57 \times 10^{11} \text{ atoms of } ^{14}\text{C} \\
 \text{Decay constant } = r &= \frac{\ln 2}{t_{1/2}} = \frac{\ln 2}{5730 \text{ y}} = 2.3 \times 10^{-10} \text{ min}^{-1} \\
 \text{Rate of decay} &= R_0 = rN_0 = 105 \text{ decays/min}
 \end{aligned}$$

(b) If $n = \#$ of half-lives:

$$\begin{aligned}
 R &= 10 \text{ decays/min} = \frac{R_0}{2^n} \\
 2^n &= \frac{R_0}{R} \Rightarrow n \log 2 = \log \frac{R_0}{R} \Rightarrow n = \frac{\log (R_0/R)}{\log 2} = \frac{\log 10.5}{\log 2} = 3.39 \text{ half-lives} = 3.39 t_{1/2} \\
 3.39 t_{1/2} &= 3.39 (5730 \text{ years}) = 19,400 \text{ years.}
 \end{aligned}$$

2. β Decay T & Z #13.19. $^{14}\text{C} \rightarrow ^{14}\text{N} + \bar{e} + \bar{\nu}$.

$$\begin{aligned}
 K &= \Delta Mc^2 = [m_{atom}(^{14}\text{C}) - m_{atom}(^{14}\text{N})] c^2 = [14.003242 - 14.003074] uc^2 \\
 &= 0.000168 (931.5 \text{ MeV}/c^2) c^2 = 0.156 \text{ MeV.}
 \end{aligned}$$

3. **Radioactive Series.** T & Z #13.27.

For a radioactive decay to occur, the total mass after the decay must be less than the total mass of the parent objects. In other words, kinetic energy must be positive.

<i>reaction</i>	<i>comment</i>	<i>possible?</i>
$^{228}\text{Th} \rightarrow ^{227}\text{Th} + n$	$K = \Delta Mc^2 = (M_i - M_f)c^2 = -7.13 \text{ MeV}$	No
$\rightarrow ^{227}\text{Ac} + p$	$K = -6.39 \text{ MeV}$	No
$\rightarrow ^{228}\text{Pa} + \bar{\nu} + e^-$	$K = -2.10 \text{ MeV}$	No
$\rightarrow ^{224}\text{Ra} + \alpha$	$K = 5.50 \text{ MeV}$	Yes
$\rightarrow ^{216}\text{Po} + ^{12}\text{C}$	$K = 24.99 \text{ MeV}$	Yes




Figure 1: $X \rightarrow Y + \alpha$.

4. **Radioactive Series.** T & Z #13.34.

(a) By conservation of momentum, $|p_\alpha| = |p_Y| \equiv p$. Because we can treat these particles nonrelativistically, $K = p^2/2m$ which means that $K_\alpha = p^2/2m_\alpha = p^2/2m_\alpha$ and $K_Y = p^2/2m_Y = p^2/2m_Y$. Taking the ratio of kinetic energies we see:

$$\frac{K_\alpha}{K_Y} = \frac{m_Y}{m_\alpha} \rightarrow K_\alpha = \left(\frac{m_Y}{m_\alpha}\right) K_Y$$

(b) Typical $m_Y \approx 220$ u, $m_\alpha = 4 \rightarrow K_\alpha = 55K_Y = (55/56)K_{\text{total}} \approx 98\%K_{\text{total}}$.

(c) $X \rightarrow Y + e$. $m_e = 5 \times 10^{-4}$ u, $m_Y = 1$ u to 240u. Therefore, $K_e = (m_Y/m_e)K_Y \rightarrow 99.95\%$ to 99.9998% of K_{total} .

Actually these are maximum values in (b) and (c) because some of the kinetic energy will go to a neutrino.

5. T&Z #13.38.




Figure 2:

(a) $R = R_0 a^{1/3} \rightarrow R(^1H) = 1.07$ fm, $R(^4He) = 1.70$ fm. d = distance between nuclear centers = $1 + 1.07 + 1.70$ fm = 3.77 fm. Energy needed to bring one of the particles from infinity is just the potential energy:

$$E = \frac{kq_1q_2}{d} = \frac{2ke^2}{d} = \frac{2(1.44 \text{ MeV fm})}{3.77 \text{ fm}} = 0.76 \text{ MeV}.$$

(b) $R(^{238}U) = 6.63$ fm $\rightarrow d = 1 + 1.70 + 6.63 = 9.33$ fm.

$$E = \frac{kq_1q_2}{d} = \frac{2(92)ke^2}{d} = \frac{184(1.44 \text{ MeV fm})}{9.33 \text{ fm}} = 28 \text{ MeV}.$$

6. T & Z, #13.43.

(a) $^{207}_{84}\text{Po} \rightarrow ^{207}_{85}\text{At} + e^- + \bar{\nu}$. Nucleon number and charge are conserved, but $\Delta K = (M_i + M_f)c^2 < 0$. Not possible.

(b) ${}_{86}^{213}\text{Rn} \rightarrow {}_{84}^{208}\text{Po} + {}_2^4\text{He}$. Nucleon number is not conserved. Not possible.

(c) ${}_{84}^{212}\text{Po} \rightarrow {}_{82}^{208}\text{Pb} + {}_2^4\text{He}$. Nucleon number and charge are conserved, and $\Delta K > 0$. Possible.

(d) ${}_1^1\text{H} + {}_7^{14}\text{N} \rightarrow {}_6^{12}\text{C} + {}_2^4\text{He}$. Nucleon number is not conserved. Not possible.

(e) $n + {}_{11}^{22}\text{Na} \rightarrow {}_{10}^{22}\text{Ne} + {}_1^1\text{H}$. Nucleon number and charge are conserved, and $\Delta K > 0$. Possible.